

The Space of Local Fields as a Module over the Ring of Local Integrals of Motion

Atsushi Nakayashiki*

Abstract

The chiral space of local fields in Sine-Gordon or the SU(2)-invariant Thirring model is studied as a module over the commutative algebra $\mathcal{D}$ of local integrals of motion. Using the recent construction of form factors by means of quantum affine algebra at root of unity due to Feigin et al. we construct a $\mathcal{D}$ -free resolution of the space of local fields. In general the cohomologies of the de Rham type complex associated with the space of local fields are determined and shown to be the irreducible representations of the symplectic group $Sp(2\infty)$. Babelon-Bernard-Smirnov's description of the space of local fields automatically incorporated in this framework.

1 Introduction

Let $U_{\sqrt{-1}}(\widehat{sl_2})$ be the quantum affine algebra at $q = \sqrt{-1}$, $x_k^\pm$, b_n (a part of) Drinfeld generators, $V_{\sqrt{-1}}(\Lambda_i)$ the level one representation and $V_{\sqrt{-1}}(\Lambda_i)_{2m+i}$ the subspace of vectors whose sl_2 weight is $2m+i$.

In [4] Feigin et al. proved that the chiral subspace of local fields in the quantum sine-Gordon model at generic coupling constant and that in the SU(2) invariant Thirring model (ITM) are isomorphic to the space

$$A_{2m+i} = \frac{V_{\sqrt{-1}}(\Lambda_i)_{2m+i}}{x_0^- V_{\sqrt{-1}}(\Lambda_i)_{2m+i+2} + (x_0^-)^{(2)} V_{\sqrt{-1}}(\Lambda_i)_{2m+i+4}}, \quad m \geq 0,$$

where $(x_0^-)^{(2)} = (x_0^-)^2 / (q + q^{-1})$ is the divided power of x_0^- . The local integral of motion with spin s , s being odd, is identified with b_s . They become central at $q = \sqrt{-1}$ and A_{2m+i} becomes a module over the commutative ring

$$\mathcal{D} = \mathbb{C}[b_{-1}, b_{-3}, \dots].$$

To study the structure of A_{2m+i} as a $\mathcal{D}$ -module is an interesting problem because it is a common problem for any integrable system. In particular it will give a suitable framework to compare structures of different integrable systems such as massive integrable field theories, conformal field theories and classical integrable systems [1, 14].

The main aim of this article is to determine the $\mathcal{D}$ -module structure of A_{2m+i} completely. More precisely we construct a $\mathcal{D}$ -free resolution of A_{2m+i} . Each term of the resolution is described in terms of the Fock space of free fermions.

*Faculty of Mathematics, Kyushu University, Ropponmatsu 4-2-1, Fukuoka 810-8560, Japan, e-mail: 6vertex@math.kyushu-u.ac.jp

The problem to study the space of local fields as a module over the ring of local integrals of motion was initiated by Babelon-Bernard-Smirnov [1]. They studied the restricted sine-Gordon model and derived the $c = 1$ character of the Virasoro algebra. The important observation in [1] is that the fermionization simplifies the description of polynomials giving rise to null vectors. The description of the space of local fields in terms of the representations of the quantum affine algebra automatically incorporates the fermions and related operators in [1]. We remark here that it is difficult to determine the $\mathcal{D}$ -module structure of A_{2m+i} in the previous description by minimal form factors [10, 11, 5, 6].

The commutative ring $\mathcal{D}$ and a module over it determines a de Rham type complex. Let

$$\Omega^{\frac{\infty}{2}} = \mathbb{C} dt_{-1} \wedge dt_{-3} \wedge \cdots,$$

be the space of highest degree forms and $\Omega^{\frac{\infty}{2}-p}$ be the vector space generated by differential forms which are obtained from $dt_{-1} \wedge dt_{-3} \wedge \cdots$ by removing p dt_i 's. Then the pair of the vector space and a differential

$$C_{2m+i}^{\frac{\infty}{2}-p} = A_{2m+i} \otimes \Omega^{\frac{\infty}{2}-p}, \quad (1)$$

$$d = \sum_{s=1}^{\infty} b_{-(2s-1)} \otimes dt_{-(2s-1)}, \quad (2)$$

defines a complex. The highest cohomology group is isomorphic to

$$\frac{A_{2m+i}}{\sum_{s=1}^{\infty} b_{-(2s-1)} A_{2m+i}}.$$

This space is the space of chiral local fields modulo the action of local integrals of motion and its basis gives the minimal set of generators of A_{2m+i} as a $\mathcal{D}$ -module. Thus to determine it is in fact a first step to construct a free resolution of A_{2m+i} . In turn, using the free resolution of A_{2m+i} it is possible to describe all cohomology groups in terms of the Fock space of free fermions. We show that those cohomology groups become the irreducible representations of the symplectic group $Sp(2\infty)$, which is obtained as the inductive limit of $Sp(2n)$'s. This result suggests that the cohomology groups of (1), (2) give the universal structure of the cohomology groups of affine hyperelliptic Jacobian varieties [9, 12]. Thus one can think of the result as an example of the comparison of different integrable systems mentioned above. We shall study this subject in a subsequent paper.

The present paper is organized in the following manner. After introduction the notations concerning the quantum affine algebra at $q = \sqrt{-1}$ and its level one representations are explained in section two. Then the chiral space of local fields are defined using them. In section 3 fermions are introduced and the integral expressions of x_0^- and $(x_0^-)^{(2)}$ are given in terms of them. The free resolution of A_{2m} are constructed in section 4. In section five the de Rham type cohomology groups associated with A_{2m} are determined. The space A_{2m+1} is studied in section 6. We prove that A_{2m+1} is a free $\mathcal{D}$ -module. In section 7 the cohomology groups are shown to be the irreducible representations of the symplectic group.

2 The chiral space

We recall the results of [4]. Notations mainly follows that paper. Let q be an indeterminate. The quantum affine algebra $U_q(\widehat{sl_2})$ is the $\mathbb{C}(q)$ Hopf algebra generated by $x_k^{\pm}$ ($k \in \mathbb{Z}$), b_n ($n \in \mathbb{Z} \setminus \{0\}$), $t_1^{\pm 1}$, $C^{\pm 1}$, $D^{\pm}$ with the following defining relations:

$$\begin{aligned}
& C^{\pm 1} \text{ is central, } CC^{-1} = C^{-1}C = 1, \\
& t_1 t_1^{-1} = t_1^{-1} t_1 = 1, \quad t_1 b_n t_1^{-1} = b_n, \quad t_1 x_n^{\pm} t_1^{-1} = q^{\pm 2} x_n^{\pm}, \\
& DD^{-1} = D^{-1}D = 1, \quad Dt_1 = t_1 D, \\
& Db_n D^{-1} = q^n b_n, \quad Dx_n^{\pm} D^{-1} = q^n x_n^{\pm}, \\
& [b_m, b_n] = m \frac{[2m]}{[m]^2} \frac{C^m - C^{-m}}{q - q^{-1}} \delta_{m+n,0}, \\
& [b_n, x_k^{\pm}] = \pm \frac{[2n]}{[n]} C^{\frac{n \mp |n|}{2}} x_{k+n}^{\pm}, \\
& x_{k+1}^{\pm} x_l^{\pm} - q^{\pm 2} x_l^{\pm} x_{k+1}^{\pm} = q^{\pm 2} x_k^{\pm} x_{l+1}^{\pm} - x_{l+1}^{\pm} x_k^{\pm}, \\
& [x_k^+, x_l^-] = \frac{C^{-l} \varphi_{k+l}^+ - C^{-k} \varphi_{k+l}^-}{q - q^{-1}},
\end{aligned}$$

where

$$\sum_{k \in \mathbb{Z}} \varphi_{\pm k}^{\pm} z^k = t_1^{\pm 1} \exp \left(\pm \sum_{n=1}^{\infty} \frac{q^n - q^{-n}}{n} \right) b_{\pm n} z^n.$$

Since we do not use the coproduct in this paper we omit the description it.

In order to specialize q to a complex number one has to define the so called integral form. To this end we need some notations. For $x \in U_q(\widehat{sl_2})$, the divided power of x is defined by

$$x^{(r)} = \frac{x^r}{[r]!}, \quad [r] = \frac{q^r - q^{-r}}{q - q^{-1}}, \quad [r]! = [r] \cdots [2][1].$$

Let $A = \mathbb{C}[q, q^{-1}]$. Then the integral form U_A is defined to be the A -subalgebra of $U_q(\widehat{sl_2})$ generated by $(x_k^{\pm})^{(r)}$ ($k \in \mathbb{Z}, r \geq 0$), b_n $n \in \mathbb{Z} \setminus \{0\}$, $t_1^{\pm 1}$, $C^{\pm 1}$, $D^{\pm 1}$.

The algebra $U_{\sqrt{-1}}(\widehat{sl_2})$ is then defined by

$$U_{\sqrt{-1}}(\widehat{sl_2}) = \left(\frac{A}{(q - \sqrt{-1})A} \right) \otimes_A U_A.$$

We denote the subalgebra of U_A without $D^{\pm 1}$ by U'_A and similarly for $U'_{\sqrt{-1}}(\widehat{sl_2})$.

There are two notable properties which distinguish $U'_{\sqrt{-1}}(\widehat{sl_2})$ from the generic q case. They are given by the equations

$$x_k^{\pm} x_l^{\pm} = -x_l^{\pm} x_k^{\pm}, \quad (3)$$

$$[b_{2n-1}, U'_{\sqrt{-1}}(\widehat{sl_2})] = 0 \quad \forall n \in \mathbb{Z}. \quad (4)$$

Let α_0, α_1 and Λ_0, Λ_1 be the simple roots and the fundamental weights of $\widehat{sl_2}$ respectively. The Frenkel-Jing realization of the level one integrable highest weight representation $V(\Lambda_i)$ of $U_q(\widehat{sl_2})$ determines the level one representation of $U_{\sqrt{-1}}(\widehat{sl_2})$ [2]. Define the integral form $V_A(\Lambda_i)$ by the free A -module

$$V_A(\Lambda_i) = \oplus_{m \in \mathbb{Z}} A[b_n \mid n < 0] e^{\Lambda_i + m\alpha_1}.$$

The action of U_A on this space is defined in the following way. For $Pe^\beta \in V_A(\Lambda_i)$ define the actions of C, ∂ by

$$\begin{aligned} C(Pe^\beta) &= q(Pe^\beta), \\ \partial(Pe^\beta) &= \langle h_1, \beta \rangle (Pe^\beta), \end{aligned}$$

where $\langle h_1, \alpha_1 \rangle = 2$, $\langle h_1, \Lambda_j \rangle = \delta_{1,j}$. For $n < 0$ b_n acts by the multiplication by itself and for $n > 0$ it acts by taking the commutator $[b_n, \cdot]$. Define the grading on $V_A(\Lambda_i)$ as

$$\deg b_n = n, \quad \deg e^{\Lambda_i + m\alpha_1} = -m^2 - im.$$

Let

$$x^\pm(z) = \sum_{n \in \mathbb{Z}} x_n^\pm z^{-n-1}.$$

Then the generators of U_A acts on $V_A(\Lambda_i)$ by the following formulas

$$\begin{aligned} x^+(z) &= \exp\left(\sum_{n=1}^{\infty} \frac{b_{-n}}{n} z^n\right) \exp\left(-\sum_{n=1}^{\infty} \frac{b_n}{n} (qz)^{-n}\right) e^{\alpha_1} z^\partial, \\ x^-(z) &= \exp\left(-\sum_{n=1}^{\infty} \frac{b_{-n}}{n} (qz)^n\right) \exp\left(\sum_{n=1}^{\infty} \frac{b_n}{n} z^{-n}\right) e^{-\alpha_1} z^{-\partial}, \\ t_1 &= q^\partial, \quad Du = q^{\deg u} u \quad \text{for a homogeneous } u \in V_A(\Lambda_i). \end{aligned} \tag{5}$$

We set

$$V_A(\Lambda_i)_{2m+i} = A[b_n \mid n < 0] e^{\Lambda_i + m\alpha_1}.$$

Then the space

$$V_{\sqrt{-1}}(\Lambda_i) = \left(\frac{A}{(q - \sqrt{-1})A}\right) \otimes_A V_A(\Lambda_i),$$

becomes a $U_{\sqrt{-1}}(\widehat{sl_2})$ -module and

$$V_{\sqrt{-1}}(\Lambda_i)_{2m+i} = \left(\frac{A}{(q - \sqrt{-1})A}\right) \otimes_A V_A(\Lambda_i)_{2m+i},$$

becomes its subspace.

Set

$$A_{2m+i} = \frac{V_{\sqrt{-1}}(\Lambda_i)_{2m+i}}{x_0^- V_{\sqrt{-1}}(\Lambda_i)_{2m+i+2} + (x_0^-)^{(2)} V_{\sqrt{-1}}(\Lambda_i)_{2m+i+4}}, \quad i = 0, 1, \quad m \geq 0.$$

Because of (4) it becomes a module over the commutative algebra

$$\mathcal{D} = \mathbb{C}[b_{-1}, b_{-3}, \dots].$$

The chiral space of local fields in SG-model at generic coupling constant and that in $SU(2)$ -ITM are both isomorphic to A_{2m+i} [4, 5, 6]. In SG case the fields are highest weight vectors with the weight $2m+i$ with respect to a certain quantum group $U_p(\widehat{sl_2})$ for some p and in $SU(2)$ -ITM case the fields are highest weight vectors with weight $2m+i$ respect to sl_2 . Local integrals of motion with spin $2s-1$ are identified with b_{2s-1} . Since we consider the chiral space, only the local integrals of motion with negative spins act on it.

3 Fermions

We consider the Neveu-Schwarz(NS) and Ramond(R) fermions $\{\psi_{2n-1}, \psi_{2n-1}^* \mid n \in \mathbb{Z}\}$ and $\{\psi_{2n}, \psi_{2n}^* \mid n \in \mathbb{Z}\}$ respectively. They satisfy the canonical anti-commutation relations

$$[\psi_m, \psi_n]_+ = [\psi_m^*, \psi_n^*]_+ = 0, \quad [\psi_m, \psi_n^*]_+ = \delta_{m,n}.$$

In the following, objects with odd indices are for NS-fermions and those with even indices are for R-fermions. The vacuum vectors $|m\rangle$ and $\langle m|$ are introduced by the following relations

$$\begin{aligned} \langle m|\psi_n &= 0 \quad \text{for } n \leq m, & \langle m|\psi_n^* &= 0 \quad \text{for } n > m, \\ \psi_n|m\rangle &= 0 \quad \text{for } n > m, & \psi_n^*|m\rangle &= 0 \quad \text{for } n \leq m. \end{aligned}$$

These vacuums are related by

$$\psi_m^*|m-2\rangle = |m\rangle, \quad \langle m-2|\psi_m = \langle m|.$$

The Fock spaces H_m, H_m^* are constructed from $|m\rangle$ and $\langle m|$ respectively by the same number of ψ_k and ψ_l^* . The pairing between H_m and H_m^* are defined by the condition

$$\langle m|m\rangle = 1.$$

The fermion operators are introduced as

$$\begin{aligned} \psi(z) &= \sum_{n \in \mathbb{Z}} \psi_{2n+1} z^{-2n-1}, \quad \psi^*(z) = \sum_{n \in \mathbb{Z}} \psi_{2n+1}^* z^{2n+1} \quad \text{for NS-fermions,} \\ \psi(z) &= \sum_{n \in \mathbb{Z}} \psi_{2n} z^{-2n}, \quad \psi^*(z) = \sum_{n \in \mathbb{Z}} \psi_{2n}^* z^{2n} \quad \text{for R-fermions.} \end{aligned}$$

Let $\tilde{b}_{2n}$ ($n \in \mathbb{Z}$) satisfy the canonical commutation relations

$$[\tilde{b}_{2m}, \tilde{b}_{2n}] = \delta_{m+n,0}. \quad (6)$$

We set

$$H(b) = \sum_{l=1}^{\infty} \tilde{b}_{-2l} h_{2l}, \quad h_{2l} = \sum_n \psi_n \psi_{n-2l}^*,$$

where the summation in n of h_{2l} is taken over odd $n \in \mathbb{Z}$ for NS-fermions and even n for R-fermions.

The boson-fermion correspondence gives the isomorphism of Fock spaces [3]

$$\begin{aligned} H_{2m+i} &\simeq \mathbb{C}[\tilde{b}_{-2}, \tilde{b}_{-4}, \dots] e^{\Lambda_i + m\alpha_1}, \\ a|2m+i\rangle &\mapsto \langle 2m+i| e^{H(b)} a|2m+i\rangle. \end{aligned}$$

By this correspondence the fermion operators are written as [3]

$$\begin{aligned} \psi(z) &= \exp\left(\sum_{n=1}^{\infty} \tilde{b}_{-2n} z^{2n}\right) \exp\left(-\sum_{n=1}^{\infty} \frac{\tilde{b}_{2n}}{n} z^{-2n}\right) e^{-\alpha} z^{-\partial}, \\ \psi^*(z) &= \exp\left(-\sum_{n=1}^{\infty} \tilde{b}_{-2n} z^{2n}\right) \exp\left(\sum_{n=1}^{\infty} \frac{\tilde{b}_{2n}}{n} z^{-2n}\right) z^{\partial} e^{\alpha} \end{aligned}$$

for both NS and R.

Two bosons b_{2n} and $\tilde{b}_{2n}$ are related by

$$\tilde{b}_{2n} = \begin{cases} \frac{-1}{2^n} b_{2n}, & n \geq 1 \\ \frac{(-1)^n}{2n} b_{2n}, & n \leq -1. \end{cases}$$

Then we have

Proposition 1 *By the boson-fermion correspondence x_0^- and $(x_0^-)^{(2)}$ on $V(\Lambda_i)$ are expressed as*

$$x_0^- = \int \frac{dz}{2\pi i} e^{X(z)} \psi(z), \quad (7)$$

$$(x_0^-)^{(2)} = \frac{-i}{2} \int \int_{|z_1| > |z_2|} \frac{dz_1}{2\pi i} \frac{dz_2}{2\pi i} \tau\left(\frac{z_2}{z_1}\right) e^{X(z_1)} e^{X(z_2)} \psi(z_1) \psi(z_2), \quad (8)$$

where

$$X(z) = \sum_{n=1}^{\infty} \frac{\tilde{b}_{-(2n-1)}}{2n-1} z^{2n-1},$$

$$\tau(z) = \sum_{n=1}^{\infty} z^{2n-1} - 2 \sum_{n=1}^{\infty} z^{2n},$$

and $\psi(z)$ is R-fermion for $i = 0$ and NS-fermion for $i = 1$.

Remark. The right hand sides of (7) and (8) are similar to the operators appeared in [1]. The only difference, besides the overall constant multiples, is the coefficients of even and odd powers of z in $\tau(z)$.

Proof. The expression for x_0^- follows from (4) and the definitions of $\psi(z)$, $X(z)$. Let us prove (8). For $n \geq 1$, b_n can be written as

$$b_n = n(q^n + q^{-n}) \frac{\partial}{\partial b_{-n}}.$$

We substitute this into (5). In this description of $x^-(z)$ we have

$$(x_0^-)^{(2)} = \frac{1}{2} \lim_{q \rightarrow i} \frac{(x_0^-)^2}{q - i} = \frac{1}{2} \int \int \frac{dz_1}{2\pi i} \frac{dz_2}{2\pi i} \frac{d}{dq} \left(x^-(z_1; q) x^-(z_2; q) \right) \Big|_{q=i},$$

where we write the q -dependence of $x^-(z)$ explicitly and denote $i = \sqrt{-1}$. By calculation

$$\begin{aligned} & \left(\frac{d}{dq} x^-(z; q) \right) \Big|_{q=i} \\ &= i \sum_{n=1}^{\infty} b_{-n} i^n z^n x^-(z_1; i) + x^-(z; q) \sum_{n=1}^{\infty} (-1)^{n-1} (2n-1) z^{-(2n-1)} \frac{\partial}{\partial b_{-(2n-1)}}. \end{aligned}$$

Then

$$\begin{aligned}
& \frac{d}{dq} \left(x^-(z_1; q) x^-(z_2; q) \right) |_{q=i} \\
&= i \sum_{n=1}^{\infty} b_{-n} i^n (z_1^n + z_2^n) x^-(z_1; i) x^-(z_2; i) \\
&+ x^-(z_1; i) x^-(z_2; i) \sum_{n=1}^{\infty} (-1)^{n-1} (2n-1) (z_1^{-(2n-1)} + z_2^{-(2n-1)}) \frac{\partial}{\partial b_{-(2n-1)}} \\
&- i \tau \left(\frac{z_2}{z_1} \right) x^-(z_1; i) x^-(z_2; i).
\end{aligned}$$

Here we use (4) and the commutation relations of b_n . Due to (3)

$$\int \int \frac{dz_1}{2\pi i} \frac{dz_2}{2\pi i} (z_1^n + z_2^n) x^-(z_1; i) x^-(z_2; i) = 0 \quad \forall n \in \mathbb{Z}.$$

■

4 Resolution of A_{2m}

In terms of the Fock space of fermions A_{2m} is written as

$$A_{2m} \simeq \frac{\mathcal{D} \otimes H_{2m}}{x_0^- (\mathcal{D} \otimes H_{2m+2}) + (x_0^-)^{(2)} (\mathcal{D} \otimes H_{2m+4})}.$$

Since $(x_0^-)^2 = 0$ and $[x_0^-, (x_0^-)^{(2)}] = 0$ it is possible to define a complex

$$\cdots \longrightarrow \frac{\mathcal{D} \otimes H_{2m+2}}{(x_0^-)^{(2)} (\mathcal{D} \otimes H_{2m+6})} \longrightarrow \frac{\mathcal{D} \otimes H_{2m}}{(x_0^-)^{(2)} (\mathcal{D} \otimes H_{2m+4})} \longrightarrow A_{2m} \longrightarrow 0, \quad (9)$$

where the map

$$\frac{\mathcal{D} \otimes H_{2n}}{(x_0^-)^{(2)} (\mathcal{D} \otimes H_{2n+4})} \longrightarrow \frac{\mathcal{D} \otimes H_{2n-2}}{(x_0^-)^{(2)} (\mathcal{D} \otimes H_{2n+2})}$$

is given by the left multiplication by x_0^- .

Proposition 2 *The module*

$$\frac{\mathcal{D} \otimes H_{2n}}{(x_0^-)^{(2)} (\mathcal{D} \otimes H_{2n+4})}$$

is a free $\mathcal{D}$ -module.

Proof. To prove the proposition we write x_0^- and $(x_0^-)^{(2)}$ in the component form. Let us write

$$e^{X(z)} = \sum_{n=0}^{\infty} T_{-n} z^n.$$

Notice that $T_{-(2n-1)}$ is a homogeneous polynomial of $\tilde{b}_{-(2m-1)}$ with the degree $-(2n-1)$ and it has the form

$$T_{-(2n-1)} = \frac{\tilde{b}_{-(2n-1)}}{2n-1} + \dots,$$

where $\dots$ part does not contain $\tilde{b}_{-(2n-1)}$. In particular

$$\mathcal{D} = \mathbb{C}[T_{-1}, T_{-3}, \dots].$$

By calculation we have

$$\begin{aligned} x_0^- &= \sum_{n=1}^{\infty} T_{-(2n-1)} \psi_{2n}, \\ (x_0^-)^{(2)} &= \frac{-i}{2} \sum_{n=1}^{\infty} \left(\psi_{2(-n+1)} + \sum_{l=1}^{\infty} Q_{n-1,l}(T) \psi_{2(-n+1+l)} \right) \psi_{2n}, \end{aligned}$$

where

$$Q_{n,l}(T) = \sum_{n_1+n_2=l, 0 \leq n_1, 0 \leq n_2 \leq n} (T_{-2n_1} T_{-2n_2} - 2T_{-(2n_1+1)} T_{-(2n_2-1)}).$$

We omit the tensor symbol $\otimes$ in writing elements of $\mathcal{D} \otimes H_{2m}$ etc. for the sake of simplicity. Using this expression we make a base change of $\mathcal{D} \otimes H_{2n}$. Let us define $\{\tilde{\psi}_{2n}, \tilde{\psi}_{2n}^*\}$ in the following way. First we set

$$\begin{aligned} \tilde{\psi}_{2n} &= \psi_{2n} \quad \text{for } n \geq 1, \\ \tilde{\psi}_{-2n} &= \psi_{-2n} + \sum_{l=1}^{\infty} Q_{n,l}(T) \psi_{2(-n+l)} \quad \text{for } n \geq 0, \end{aligned}$$

and write it as

$$\tilde{\psi}_i = \sum_j B_{ij} \psi_j,$$

where i, j are even integers. Then the matrix $B = (B_{ij})$ is a triangular matrix such that its diagonal entries are all 1 and $B_{ij} = \delta_{ij}$ for $i \geq 2$. Thus B^{-1} exists. Define the matrix C by

$$C = (c_{ij}) = {}^t(B^{-1})$$

and set

$$\tilde{\psi}_i^* = \sum_j c_{ij} \psi_j^*.$$

Then $\{\tilde{\psi}_k, \tilde{\psi}_l^*\}$ satisfy the canonical anti-commutation relations

$$[\tilde{\psi}_k, \tilde{\psi}_l]_+ = [\tilde{\psi}_k^*, \tilde{\psi}_l^*]_+ = 0, \quad [\tilde{\psi}_k, \tilde{\psi}_l^*]_+ = \delta_{k,l}.$$

Moreover the vacuum is the same for $\{\tilde{\psi}_k, \tilde{\psi}_l^*\}$, that is, the following relations hold

$$\begin{aligned} \tilde{\psi}_{2n} |2m\rangle &= 0 \quad \text{for } n > m, & \tilde{\psi}_{2n}^* |2m\rangle &= 0 \quad \text{for } n \leq m, \\ \langle 2m | \tilde{\psi}_{2n} &= 0 \quad \text{for } n \leq m, & \langle 2m | \tilde{\psi}_{2n}^* &= 0 \quad \text{for } n > m. \end{aligned}$$

Therefore we denote the vacuum for $\{\tilde{\psi}_k, \tilde{\psi}_l^*\}$ by the same symbol as for $\{\psi_k, \psi_l^*\}$. We denote the Fock space of $\{\tilde{\psi}_k, \tilde{\psi}_l^*\}$ by $\tilde{H}_{2m}, \tilde{H}_{2m}^*$. Then we have the isomorphism of $\mathcal{D}$ modules

$$\mathcal{D} \otimes H_{2m} \simeq \mathcal{D} \otimes \tilde{H}_{2m}.$$

In terms of $\{\tilde{\psi}_k, \tilde{\psi}_l^*\}, x_0^-$ and $(x_0^-)^{(2)}$ take simple forms

$$\begin{aligned} x_0^- &= \sum_{n=1}^{\infty} T_{-(2n-1)} \tilde{\psi}_{2n}, \\ (x_0^-)^{(2)} &= \frac{i}{2} \sum_{n=1}^{\infty} \tilde{\psi}_{-2(n-1)} \tilde{\psi}_{2n}. \end{aligned}$$

In particular $(x_0^-)^{(2)}$ defines a map

$$(x_0^-)^{(2)} : \tilde{H}_{2n} \longrightarrow \tilde{H}_{2n-4}. \quad (10)$$

Thus we have the isomorphism

$$\frac{\mathcal{D} \otimes H_{2n}}{(x_0^-)^{(2)}(\mathcal{D} \otimes H_{2n+4})} \simeq \mathcal{D} \otimes \frac{\tilde{H}_{2n}}{(x_0^-)^{(2)}\tilde{H}_{2n+4}}.$$

This proves the proposition. ■

Remark. The argument used in the proof of the proposition is essentially due to Babelon-Bernard-Smirnov[1].

We set

$$W_{2n} = \frac{\tilde{H}_{2n}}{(x_0^-)^{(2)}\tilde{H}_{2n+4}}.$$

Theorem 1 *The complex (9)*

$$\cdots \longrightarrow \mathcal{D} \otimes W_{2m+2} \longrightarrow \mathcal{D} \otimes W_{2m} \longrightarrow A_{2m} \longrightarrow 0,$$

is a $\mathcal{D}$ -free resolution of A_{2m} .

The following lemmas are sufficient to prove the theorem.

Lemma 1 *The complex*

$$\cdots \longrightarrow \mathcal{D} \otimes \tilde{H}_4 \longrightarrow \mathcal{D} \otimes \tilde{H}_2 \longrightarrow \mathcal{D} \otimes \tilde{H}_0 \longrightarrow 0$$

is exact at $\tilde{H}_{2n}$ $n \geq 1$, where the maps are defined by the multiplication by x_0^- .

Lemma 2 *The map (10) is injective for $n \geq 1$.*

Assuming these lemmas we give a proof of the theorem first.

Proof of Theorem 1.

It is sufficient to prove the exactness at $\mathcal{D} \otimes W_{2n}$, $n > m$. Suppose that $v \in \tilde{H}_{2n}$ satisfies

$$x_0^- v = (x_0^-)^{(2)} w$$

for some $w \in \tilde{H}_{2n+2}$. Then

$$(x_0^-)^{(2)}(x_0^- w) = 0.$$

Since $(x_0^-)^{(2)}$ is injective by Lemma 2, $x_0^- w = 0$. By Lemma 1, $w = x_0^- u$ for some $u \in \tilde{H}_{2n+4}$. Then

$$x_0^-(v - (x_0^-)^{(2)}u) = 0,$$

and again by Lemma 1

$$v - (x_0^-)^{(2)}u = x_0^- y$$

for some $y \in \tilde{H}_{n+2}$. This proves the theorem. ■

Proof of Lemma 1.

Notice that

$$\tilde{H}_{2m} = \sum \mathbb{C} \tilde{\psi}_{2r_1}^* \cdots \tilde{\psi}_{2r_{k+m}}^* \tilde{\psi}_{2s_1} \cdots \tilde{\psi}_{2s_k} |0\rangle,$$

where the summation is taken for all

$$0 < r_1 < \cdots < r_{k+m}, \quad s_1 < \cdots < s_k \leq 0. \quad (11)$$

We set

$$d = x_0^- = \sum_{s=1}^{\infty} T_{-(2s-1)} \psi_{2s},$$

for the sake of simplicity. For each $(s_1, \dots, s_k)$ satisfying (11) we set

$$\tilde{H}_{2m}(s_1, \dots, s_k) = \sum_{0 < r_1 < \cdots < r_{k+m}} \mathbb{C} \tilde{\psi}_{2r_1}^* \cdots \tilde{\psi}_{2r_{k+m}}^* \tilde{\psi}_{2s_1} \cdots \tilde{\psi}_{2s_k} |0\rangle.$$

Then $\tilde{H}_{2m}$ is a direct sum of $\tilde{H}_{2m}(s_1, \dots, s_k)$'s and

$$d(\tilde{H}_{2m}(s_1, \dots, s_k)) \subset \tilde{H}_{2m-2}(s_1, \dots, s_k).$$

Thus it is sufficient to prove the exactness of the complex

$$(\mathcal{D} \otimes \tilde{H}_{2m}(s_1, \dots, s_k), d)$$

at $m \geq 1$ for each $(s_1, \dots, s_k)$. Set, for $0 < r_1 < \cdots < r_{k+m}$,

$$e_{r_1 \dots r_{k+m}} = \tilde{\psi}_{2r_1}^* \cdots \tilde{\psi}_{2r_{k+m}}^* \tilde{\psi}_{2s_1} \cdots \tilde{\psi}_{2s_k} |0\rangle.$$

They form a basis of $\tilde{H}_{2m}(s_1, \dots, s_k)$. Now suppose that

$$v \in \mathcal{D} \otimes \tilde{H}_{2m}(s_1, \dots, s_k), \quad dv = 0,$$

and write

$$v = \sum_{0 < r_1 < \cdots < r_{k+m}} P_{r_1 \dots r_{k+m}} e_{r_1 \dots r_{k+m}}, \quad P_{r_1 \dots r_{k+m}} \in \mathcal{D}.$$

If we set $t_s = T_{-(2s-1)}$ we have

$$d(e_{r_1 \dots r_{k+m}}) = \sum_{i=1}^{k+m} (-1)^{i-1} t_{r_i} e_{r_1 \dots r_{i-1}, r_{i+1} \dots r_{k+m}}.$$

Let N_1 be the maximum among $r_1, \dots, r_{k+m}$ such that $P_{r_1 \dots r_{k+m}} \neq 0$, N an integer such that $N > N_1$ and $P_{r_1 \dots r_{k+m}} \in \mathbb{C}[t_1, \dots, t_N]$ for all $P_{r_1 \dots r_{k+m}} \neq 0$. We set

$$B = \mathbb{C}[t_1, \dots, t_N], \quad K_p = \sum_{1 \leq i_1 < \dots < i_p \leq N} B e_{i_1 \dots i_p}.$$

Then

$$v \in K_{m+k}$$

and the complex

$$0 \longrightarrow K_N \xrightarrow{d} K_{N-1} \xrightarrow{d} \dots \xrightarrow{d} K_0 \longrightarrow 0$$

is the Koszul complex associated with the regular sequence $(t_1, \dots, t_N)$ of B . Thus the complex (K, d) is exact at K_n , $n \geq 1$ [8]. Since $m \geq 1$, this proves the lemma. ■

Proof of Lemma 2

Let us set

$$\omega = 2i(x_0^-)^{(2)} = - \sum_{n=1}^{\infty} \tilde{\psi}_{-2(n-1)} \tilde{\psi}_{2n}.$$

Suppose that $v \in \tilde{H}_{2m}$ satisfies

$$\omega v = 0.$$

Let us write

$$v = \sum c(r_1, \dots, r_k | s_1, \dots, s_{k+m}) \tilde{\psi}_{2r_1} \dots \tilde{\psi}_{2r_k} \tilde{\psi}_{2s_1}^* \dots \tilde{\psi}_{2s_{k+m}}^* |0\rangle,$$

where

$$r_1 < \dots < r_k \leq 0 < s_1 < \dots < s_{k+m}.$$

Let N be the maximum number among $-r_1, \dots, -r_k, s_1, \dots, s_{k+m}$ such that $c(r_1, \dots, r_k | s_1, \dots, s_{k+m}) \neq 0$. Then

$$\omega v = \omega_N v, \quad \omega_N = - \sum_{n=1}^N \tilde{\psi}_{-2(n-1)} \tilde{\psi}_{2n}.$$

Define

$$\eta_N = \sum_{n=1}^N \tilde{\psi}_{-2(n-1)}^* \tilde{\psi}_{2n}^*, \quad \xi_N = - \sum_{n=1}^N \tilde{\psi}_{-2(n-1)} \tilde{\psi}_{-2(n-1)}^* + \sum_{n=1}^N \tilde{\psi}_{2n}^* \tilde{\psi}_{2n}.$$

They act on the direct sum of the spaces

$$\tilde{H}_{2m}^{(N)} = \sum_{-N+1 < r_1 < \dots < r_k \leq 0 < s_1 < \dots < s_{k+m} \leq N} \mathbb{C} \tilde{\psi}_{2r_1} \dots \tilde{\psi}_{2r_k} \tilde{\psi}_{2s_1}^* \dots \tilde{\psi}_{2s_{k+m}}^* |0\rangle,$$

and satisfy the relations

$$[\eta_N, \omega_N] = \xi_N, \quad [\xi_N, \eta_N] = 2\eta_N, \quad [\xi_N, \omega_N] = -2\omega_N.$$

Thus ω_N , η_N and ξ_N determine the action of sl_2 on $\oplus_{m \in \mathbb{Z}} \tilde{H}_{2m}^{(N)}$ by the correspondence

$$e = \eta_N, \quad f = \omega_N, \quad h = \xi_N.$$

We have $h = m$ on $\tilde{H}_{2m}^{(N)}$. Thus by the representation theory of sl_2 , ω_N is injective at $\tilde{H}_{2m}^{(N)}$, $m \geq 1$. Since $v \in \tilde{H}_{2m}^{(N)}$, $v = 0$. ■

Let us write $D = q^d \in U_q(\widehat{sl_2})$. Then d can be considered as a degree operator of $V(\Lambda_i)$. The grading of $V(\Lambda_i)$ induces those of $V_{\sqrt{-1}}(\Lambda_i)$ and H_{2m} . Since x_0^- and $(x_0^-)^{(2)}$ are homogeneous, the quotient space A_{2m} is also graded. In general, for a graded vector space V by the degree operator d such that each homogeneous subspace is finite dimensional, we define its character by

$$\text{ch } V = \text{tr}_V(p^{-d}).$$

We remark that, with this definition of the degree, we have

$$\deg \psi_{2n} = 2n - 1, \quad \deg \psi_{2n}^* = -(2n - 1), \quad \deg |2m\rangle = -m^2.$$

Then

$$\deg \tilde{\psi}_{2n} = 2n - 1, \quad \deg \tilde{\psi}_{2n}^* = -(2n - 1),$$

and

$$\text{ch } \tilde{H}_{2m} = \text{ch } H_{2m}. \tag{12}$$

Corollary 1

$$\text{ch } A_{2m} = \frac{p^{m^2}(1 - p^{2m+1})}{(p : p)_\infty}.$$

Proof. Since

$$H_{2m} \simeq \mathbb{C}[b_{-2}, b_{-4}, \dots] e^{\Lambda_0 + m\alpha_1},$$

we have, by (12),

$$\text{ch } \tilde{H}_{2m} = \frac{p^{m^2}}{(p^2 : p^2)_\infty}.$$

By Lemma 2 ω is injective. Thus

$$\text{ch } W_{2m} = \text{ch } \tilde{H}_{2m} - \text{ch } \tilde{H}_{2m+4},$$

and

$$\text{ch } A_{2m} = \sum_{n=m}^{\infty} (-1)^{n-m} \text{ch} \left(\mathcal{D} \otimes W_{2m} \right) = \text{ch } \mathcal{D} (\text{ch } \tilde{H}_{2m} - \text{ch } \tilde{H}_{2m+2}).$$

This proves the corollary. ■

Remark. The character of A_{2m} is calculated in [10, 4] in a so called fermionic form. Combining the identity of q -series [7] with it the character formula in the corollary is proved. Here we have shown that the character formula in this bosonic form can be derived from the resolution of A_{2m} independently to other results.

Corollary 2 *We have the isomorphism*

$$\frac{A_{2m}}{\sum_{s=1}^{\infty} b_{-(2s-1)} A_{2m}} \simeq W_{2m}.$$

5 De Rham type cohomologies

Let us set

$$v_{\phi} = dt_{-1} \wedge dt_{-3} \wedge dt_{-5} \wedge \cdots.$$

For $0 > i_1 > \cdots > i_l$, i_k 's being odd, we define $v_{i_1, \dots, i_l}$ to be the formal infinite wedges obtained from v_{ϕ} by removing $dt_{i_1}, \dots, dt_{i_l}$. We define $\Omega^{\frac{\infty}{2}-p}$, $p \geq 0$ by

$$\begin{aligned} \Omega^{\frac{\infty}{2}} &= \mathbb{C} v_{\phi}, \\ \Omega^{\frac{\infty}{2}-p} &= \sum_{0 > i_1 > \cdots > i_p, i_k \text{ odd}} \mathbb{C} v_{i_1, \dots, i_p} \quad \text{for } p \geq 1 \end{aligned}$$

and set

$$C_{2m}^{\frac{\infty}{2}-p} = A_{2m} \otimes \Omega^{\frac{\infty}{2}-p}.$$

Then by defining

$$d = \sum_{s=1}^{\infty} b_{-(2s-1)} \otimes dt_{-(2s-1)},$$

to be a differential $(C_{2m}^{\frac{\infty}{2}-}, d)$ becomes a complex. We denote its cohomology group at $C_{2m}^{\frac{\infty}{2}-p}$ by $H_{2m}^{\frac{\infty}{2}-p}$. Notice that

$$H_{2m}^{\frac{\infty}{2}} \simeq \frac{A_{2m}}{\sum_{s=1}^{\infty} b_{-(2s-1)} A_{2m}},$$

which is described in Corollary 2. Other cohomology groups are described by

Proposition 3

$$H_{2m}^{\frac{\infty}{2}-p} \simeq W_{2(m+p)} \quad \text{for } p \geq 0.$$

As a consequence of this proposition we have

$$\text{ch } H_{2m}^{\frac{\infty}{2}-p} = \frac{p^{(m+p)^2}(1 - p^{4(m+p+1)})}{(p^2 : p^2)_{\infty}}.$$

Proof of Proposition 3

Consider two chain complexes of $\mathcal{D}$ -modules

$$\begin{aligned} K_p &= \mathcal{D} \otimes \Omega^{\frac{\infty}{2}-p}, & d^K &= \sum_{n=1}^{\infty} b_{-(2n-1)} \otimes dt_{-(2n-1)}, \\ L_p &= \mathcal{D} \otimes W_{2(m+p)}, & d^L &= x_0^-. \end{aligned}$$

We set $K_p = L_p = 0$ for $p < 0$. Let

$$M_{p,q} = K_p \otimes_{\mathcal{D}} L_q$$

be the double complex obtained from K and L . Here p denotes the row index and q denotes the column index. We denote by $M = (M_n)$ the total complex of $(M_{p,q})$. The bordered chain complexes of $(M_{p,q})$ in the vertical and horizontal directions are

$$H_0(M^{p,\cdot}) \simeq C_{2m}^{\frac{\infty}{2}-p}, \quad (13)$$

$$H_0(M^{\cdot,q}) \simeq W_{2(m+q)}, \quad (14)$$

respectively, where the symbol H_0 denotes the 0-th homology group of a complex. Notice that all maps in the complex (14) are zero. Therefore q -th homology group of (14) are given by

$$H_q(W_{2(m+\cdot)}) = W_{2(m+q)}.$$

By the definition p -th homology group of the complex (13) is

$$H_p(C_{2m}^{\frac{\infty}{2}-\cdot}) = H_{2m}^{\frac{\infty}{2}-p}.$$

Since two complexes

$$\begin{aligned} \cdots \longrightarrow M_{1,q} \longrightarrow M_{0,q} \longrightarrow W_{2(m+q)} \longrightarrow 0, \\ \cdots \longrightarrow M_{p,1} \longrightarrow M_{p,0} \longrightarrow C_{2m}^{\frac{\infty}{2}-p} \longrightarrow 0, \end{aligned}$$

are exact for $p, q \geq 0$, we have

$$H_n(W_{2(m+\cdot)}) \simeq H_n(M) \simeq H_n(C_{2m}^{\frac{\infty}{2}-\cdot}).$$

■

6 The structure of A_{2m+1}

The strategy is similar to the case of A_{2m} , that is, we make a base change of fermions and describe A_{2m+1} using the Fock space of new fermions. The structure of A_{2m+1} is simpler than that of A_{2m} .

The component forms of x_0^- and $(x_0^-)^{(2)}$ are given by

$$x_0^- = \sum_{n=0}^{\infty} T_{-2n} \psi_{2n+1},$$

$$(x_0^-)^{(2)} = \frac{-i}{2} \sum_{n=1}^{\infty} \left(-2\psi_{-2n+1} + \sum_{l=1}^{\infty} Q_{n-1,l}(T) \psi_{2(-n+l)+1} \right) \psi_{2n+1},$$

where

$$Q_{n,l}(T) = \sum_{n_1+n_2=l, 0 \leq n_1, 0 \leq n_2 \leq n} (-2T_{-2n_1} T_{-2n_2} + T_{-(2n_1-1)} T_{-(2n_2+1)}).$$

We set

$$\tilde{\psi}_{2n+1} = \psi_{2n+1} \quad \text{for } n \geq 0,$$

$$\tilde{\psi}_{-(2n+1)} = -2\psi_{-(2n+1)} + \sum_{l=1}^{\infty} Q_{n,l}(T) \psi_{2(-n+l)-1} \quad \text{for } n \geq 0.$$

Then we have

$$x_0^- = \sum_{n=0}^{\infty} T_{-2n} \tilde{\psi}_{2n+1},$$

$$(x_0^-)^{(2)} = \frac{-i}{2} \sum_{n=1}^{\infty} \tilde{\psi}_{-(2n-1)} \tilde{\psi}_{2n+1}.$$

We define $\tilde{\psi}_{2n+1}^*$, $n \in \mathbb{Z}$ similarly to the case of A_{2m} . Again the vacuums are invariant by this change. Thus the Fock space $\tilde{H}_{2m+1}$ is defined over the same vacuum as fermions ψ_{2n+1} , ψ_{2n+1}^* .

Let us define the subspace of $\tilde{H}_{2m+1}$ by

$$\tilde{H}_{2m+1,0} = \sum \mathbb{C} \tilde{\psi}_{2r_1+1} \cdots \tilde{\psi}_{2r_k+1} \tilde{\psi}_{2s_1+1}^* \cdots \tilde{\psi}_{2s_{k+m}+1}^* |1\rangle,$$

where $r_1 < \cdots < r_k < 0 < s_1 < \cdots < s_{k+m}$. Notice that $\tilde{\psi}_1$ is absent in $\tilde{H}_{2m+1,0}$. Since $(x_0^-)^{(2)}$ does not contain $\tilde{\psi}_1$,

$$(x_0^-)^{(2)} \tilde{H}_{2m+1,0} \subset \tilde{H}_{2m-3,0}.$$

We set

$$W_{2n+1,0} = \frac{\tilde{H}_{2n+1,0}}{(x_0^-)^{(2)} \tilde{H}_{2n+5,0}}.$$

The degree induced from $V_{\sqrt{-1}}(\Lambda_1)$ is given by

$$\deg \tilde{\psi}_{2n+1} = 2n, \quad \deg \tilde{\psi}_{2n+1}^* = -2n, \quad \deg |2m+1\rangle = -m^2 - m.$$

Theorem 2 (i) The space A_{2m+1} becomes a free $\mathcal{D}$ -module as

$$A_{2m+1} \simeq \mathcal{D} \otimes W_{2m+1,0}.$$

(ii)

$$\text{ch } A_{2m+1} = \frac{p^{m(m+1)}(1-p^{2m+2})}{(p:p)_\infty}.$$

Corollary 3

$$\frac{A_{2m+1}}{\sum_{s=1}^\infty b_{-(2s-1)} A_{2m+1}} \simeq W_{2m+1,0}.$$

Proof of Theorem 2.

Since

$$x_0^- = \tilde{\psi}_1 + T_2 \tilde{\psi}_3 + T_4 \tilde{\psi}_5 + \cdots,$$

we have the isomorphism of $\mathcal{D}$ -modules

$$\mathcal{D} \otimes \tilde{H}_{2m+1,0} \simeq \frac{\mathcal{D} \otimes \tilde{H}_{2m+1}}{x_0^- (\mathcal{D} \otimes \tilde{H}_{2m+1})}.$$

Thus the natural map

$$\mathcal{D} \otimes \tilde{H}_{2m+1,0} \longrightarrow A_{2m+1}^{(0)}$$

is surjective and its kernel is given by

$$\frac{x_0^- (\mathcal{D} \otimes \tilde{H}_{2m+1}) + (x_0^-)^{(2)} (\mathcal{D} \otimes \tilde{H}_{2m+5})}{x_0^- (\mathcal{D} \otimes \tilde{H}_{2m+1})} \simeq (x_0^-)^{(2)} (\mathcal{D} \otimes \tilde{H}_{2m+5,0}),$$

which proves (i).

Next let us prove (ii). Set

$$\begin{aligned} \omega_N &= - \sum_{n=1}^N \tilde{\psi}_{-(2n-1)} \tilde{\psi}_{2n+1}, \\ \eta_N &= \sum_{n=1}^N \tilde{\psi}_{-(2n-1)}^* \tilde{\psi}_{2n+1}^*, \\ \xi_N &= - \sum_{n=1}^N \tilde{\psi}_{-(2n-1)} \tilde{\psi}_{-(2n-1)}^* + \sum_{n=1}^N \tilde{\psi}_{2n+1}^* \tilde{\psi}_{2n+1}. \end{aligned}$$

Then they satisfy the relations of sl_2 as in the case of A_{2m} . It follows that $(x_0^-)^{(2)}$ is injective on $\tilde{H}_{2m+1,0}$, $m \geq 1$. Using the arguments of [1] we have

$$\begin{aligned} \text{ch } W_{2m+1,0} &= \text{ch } \tilde{H}_{2m+1,0} - \text{ch } \tilde{H}_{2m+5,0} \\ &= (1 - p^{2m+2}) \text{ch } \tilde{H}_{2m+1}. \end{aligned}$$

Thus (ii) follows from

$$\text{ch } \tilde{H}_{2m+1} = \frac{p^{m(m+1)}}{(p^2 : p^2)_\infty}.$$

■

We define de Rham type complex $(C_{2m+1}^{\frac{\infty}{2}-p}, d)$ and its cohomology $H_{2m+1}^{\frac{\infty}{2}-p}$ for A_{2m+1} in a similar manner to the case of A_{2m} . Then

Proposition 4

$$H_{2m+1}^{\frac{\infty}{2}-p} \simeq \begin{cases} W_{2m+1,0}, & \text{for } p = 0 \\ 0, & \text{for } p \geq 1. \end{cases}$$

The proof of this proposition is similar to that of Proposition 3 and we leave it to the reader.

7 Action of symplectic group on cohomologies

Let $Sp(2n) \subset GL(2n, \mathbb{C})$ be the symplectic group. For $n < n'$ we have the embedding

$$GL(2n, \mathbb{C}) \subset GL(2n', \mathbb{C}),$$

$$A \mapsto \begin{bmatrix} 1_{n'-n} & & \\ & A & \\ & & 1_{n'-n} \end{bmatrix},$$

where 1_r denotes the r by r unit matrix. We define the group $Sp(2\infty)$ as the inductive limit of $Sp(2n)$ with respect to this embedding. In this section we shall show that W_{2m} and $W_{2m+1,0}$ are irreducible representations of $Sp(2\infty)$.

Let us consider W_{2m} first. Set

$$\alpha_n = \tilde{\psi}_{-2(n-1)}^*, \quad \beta_n = \tilde{\psi}_{2n}^*, \quad \alpha_n^* = \tilde{\psi}_{-2(n-1)}, \quad \beta_n^* = \tilde{\psi}_{2n}, \quad n \geq 1.$$

Then

$$\omega = - \sum_{n=1}^{\infty} \alpha_n^* \beta_n^*, \quad \eta = \sum_{n=1}^{\infty} \alpha_n \beta_n. \quad (15)$$

Let

$$V_N = \oplus_{i=1}^N \mathbb{C} \alpha_i \oplus_{i=1}^N \mathbb{C} \beta_i. \quad (16)$$

We consider V_N as the vector representation of $Sp(2N)$. Then for $k \geq 1$

$$M_k^{(N)} := \frac{\wedge^k V_N}{\eta_N \wedge^{k-2} V_N}, \quad (17)$$

is isomorphic to the k -th fundamental representation of $Sp(2N)$, where

$$\eta_N = \sum_{i=1}^N \alpha_i \wedge \beta_i.$$

To each $m \geq 0$ we associate $M_{N-m}^{(N)}$, $N \geq m$. For $N < N'$ we define a map

$$\begin{aligned} \wedge^{N-m} V_N &\longrightarrow \wedge^{N'-m} V_{N'}, \\ v &\mapsto v \wedge \alpha_{N+1} \wedge \cdots \wedge \alpha_{N'}. \end{aligned} \quad (18)$$

It induces a map

$$M_{N-m}^{(N)} \longrightarrow M_{N'-m}^{(N')}. \quad (19)$$

Lemma 3 *The map (19) is injective.*

Proof. It is sufficient to prove the lemma for $N' = N + 1$. Suppose that $v \in \wedge^{N-m} V_N$ satisfy

$$v \wedge \alpha_{N+1} = \eta_{N+1} \wedge w, \quad w \in \wedge^{N-m-1} V_{N+1}. \quad (20)$$

Let us write

$$w = w' + w'' \wedge \alpha_{N+1}, \quad w' \in \wedge^{N-m-1} V_{N+1}, \quad w'' \in \wedge^{N-m-2} V_{N+1},$$

where w' and w'' do not contain α_{N+1} . Then

$$\eta_{N+1} \wedge w = \eta_N \wedge w' + (-w' \wedge \beta_{N+1} + \eta_N \wedge w'') \wedge \alpha_{N+1}.$$

By (20) we have

$$\eta_N \wedge w' = 0.$$

By the representation theory of sl_2 η_N is injective on $\wedge^{N-m-1} V_N \oplus \beta_{N+1} \wedge^{N-m-2} V_N$. Thus $w' = 0$ and we have

$$v \wedge \alpha_{N+1} = \eta_N \wedge w'' \wedge \alpha_{N+1}.$$

This shows $v = \eta_n \wedge w''$. ■

We denote $M_{\infty-m}$ the inductive limit of $M_{N-m}^{(N)}$,

$$M_{\infty-m} = \varinjlim M_{N-m}^{(N)}.$$

For $N < N'$ the subgroup $Sp(2N)$ in $Sp(2N')$ fixes $\alpha_{N+1}, \dots, \alpha_{N'}$, $\beta_{N+1}, \dots, \beta_{N'}$. Therefore $Sp(2\infty)$ acts on $M_{\infty-m}$. It is straightforward to check that this representation is irreducible.

Proposition 5

$$W_{2m} \simeq M_{\infty-m}.$$

Proof. By the representation theory of sl_2 we have the isomorphisms,

$$\begin{aligned} \omega^m : \tilde{H}_{2m} &\simeq \tilde{H}_{-2m}, \\ \omega^{m+1} \tilde{H}_{2m+4} &\simeq \eta \tilde{H}_{-2m-4}. \end{aligned}$$

Thus

$$W_{2m} = \frac{\tilde{H}_{2m}}{\omega \tilde{H}_{2m+4}} \simeq \frac{\tilde{H}_{-2m}}{\eta \tilde{H}_{-2m-4}}.$$

We shall show that W_{2m} is an inductive limit of the subspaces isomorphic to $M_{N-m}^{(N)}$.

We set, for $N \geq m$,

$$\tilde{H}_{-2m}(N) = \sum_{k=0}^{N-m} \sum \mathbb{C} \beta_{i_1} \cdots \beta_{i_k} \alpha_{j_1} \cdots \alpha_{j_{N-m-k}} | - 2N \rangle,$$

where the second summation is taken for all

$$1 \leq i_1 < \cdots < i_k \leq N, \quad 1 \leq j_1 < \cdots < j_{N-m-k} \leq N.$$

For $N < N'$ we have the inclusion

$$\begin{aligned} \tilde{H}_{-2m}(N) &\subset \tilde{H}_{-2m}(N'), \\ x | - 2N \rangle &= x \alpha_{N+1} \cdots \alpha_{N'} | - 2N' \rangle. \end{aligned} \tag{21}$$

Thus $\{\tilde{H}_{-2m}(N)\}$ defines an increasing filtration and satisfy

$$\tilde{H}_{-2m} = \cup_{N=m}^{\infty} \tilde{H}_{-2m}(N).$$

There is an isomorphism,

$$\begin{aligned} \tilde{H}_{-2m}(N) &\simeq \wedge^{N-m} V_N, \\ \beta_{i_1} \cdots \beta_{i_k} \alpha_{j_1} \cdots \alpha_{j_{N-m-k}} | - 2N \rangle &\mapsto \beta_{i_1} \wedge \cdots \wedge \alpha_{j_{N-m-k}}. \end{aligned} \tag{22}$$

It induces the isomorphism

$$\frac{\tilde{H}_{-2m}(N)}{\eta_N \tilde{H}_{-2m-4}(N)} \simeq \frac{\wedge^{N-m} V_N}{\eta_N \wedge^{N-m-2} V_N} = M_{N-m}^{(N)}.$$

By the isomorphism (22) the inclusion (21) is transformed to the map (18). Thus we have

$$\frac{\tilde{H}_{-2m}}{\eta \tilde{H}_{-2m-4}} = \lim_{\longrightarrow} \frac{\tilde{H}_{-2m}(N)}{\eta_N \tilde{H}_{-2m-4}(N)} \simeq M_{\infty-m}.$$

■

As to $W_{2m+1,0}$ we have

Proposition 6

$$W_{2m+1,0} \simeq M_{\infty-m-1}.$$

Proof. In this case we set

$$\alpha_n = \tilde{\psi}_{-(2n-1)}^*, \quad \beta_n = \tilde{\psi}_{2n+1}^*, \quad \alpha_n^* = \tilde{\psi}_{-(2n-1)}, \quad \beta_n^* = \tilde{\psi}_{2n+1}, \quad n \geq 1.$$

Then ω and η are given by (15). Define V_N and $M_k^{(N)}$ by (16) and (17) respectively. As in the case of $\tilde{H}_{2m}$ we have the isomorphisms

$$\begin{aligned} \omega^{m+1} : \tilde{H}_{2m+1,0} &\simeq \tilde{H}_{-2m-1,0}, \\ \omega^{m+2} \tilde{H}_{2m+5,0} &\simeq \eta \tilde{H}_{-2m-5,0}, \end{aligned}$$

and consequently

$$W_{2m+1,0} \simeq \frac{\tilde{H}_{-2m-1,0}}{\eta \tilde{H}_{-2m-5,0}}.$$

By the definition

$$\tilde{H}_{-2m-1,0} = \sum_{k=m+1}^{\infty} \sum \mathbb{C} \beta_{i_1} \cdots \beta_{i_{k-m-1}} \alpha_{j_1}^* \cdots \alpha_{j_k}^* |1 \rangle,$$

where the second summation is taken for all

$$1 \leq i_1 < \cdots < i_{k-m-1}, \quad 1 \leq j_1 < \cdots < j_k.$$

We identify $\tilde{H}_{-2m-1,0}$ as a subspace of $\tilde{H}_{-2m-3}$ by

$$x|1 \rangle = x\tilde{\psi}_1^*| -1 \rangle \mapsto x| -1 \rangle, \quad (23)$$

that is, by removing $\tilde{\psi}_1^*$. Since x does not contain $\tilde{\psi}_1$, this map is well-defined. Moreover the map (23) preserves the degree because $\deg \tilde{\psi}_1^* = 0$. We set

$$\tilde{H}_{-2m-1,0}(N) = \sum_{k=0}^{N-m-1} \sum \mathbb{C} \beta_{i_1} \cdots \beta_{i_k} \alpha_{j_1} \cdots \alpha_{j_{N-m-1-k}} | - (2N+1) \rangle,$$

where the second summation is taken for all

$$1 \leq i_1 < \cdots < i_k \leq N, \quad 1 \leq j_1 < \cdots < j_{N-m-1-k} \leq N.$$

Then

$$\tilde{H}_{-2m-1,0}(N) \simeq \wedge^{N-m-1} V_N. \quad (24)$$

For $N < N'$ we have the natural inclusion map

$$\begin{aligned} \tilde{H}_{-2m-1,0}(N) &\longrightarrow \tilde{H}_{-2m-1,0}(N'), \\ x| - (2N+1) \rangle &\mapsto x\alpha_{N+1} \cdots \alpha_{N'} | - (2N'+1) \rangle. \end{aligned}$$

This map is transplanted to (18) by the isomorphism (24). Thus we have

$$\frac{\tilde{H}_{-2m-1,0}}{\eta \tilde{H}_{-2m-5,0}} \simeq \lim_{\longrightarrow} \frac{\tilde{H}_{-2m-1,0}(N)}{\eta_N \tilde{H}_{-2m-5,0}(N)} \simeq M_{\infty-m-1}.$$

■

References

- [1] Babelon, O., Bernard, D. and Smirnov, F., Null-vectors in integrable field theory, *Comm. Math. Phys.* **186** (1997), 601-648.
- [2] Chari, V. and Jing, N., Realization of level one representations of $U_q(\widehat{\mathcal{G}})$ at a root of unity, *Duke Math. J* **108** (2001), 183-197.

- [3] Date, E., Jimbo, M., Kashiwara, M. and Miwa, T., Transformation groups for Soliton equations, Nonlinear Integrable Systems-Classical Theory and Quantum Theory-, Ed. by Jimbo, M. and Miwa, T., World Scientific, 1983.
- [4] Feigin, B., Jimbo, M., Kashiwara, M., Miwa, T., Mukhin, E. and Takeyama, Y., A functional model for the tensor product of level 1 highest and level -1 lowest modules for the quantum affine algebra $U_q(\widehat{sl_2})$, math.QA/0310284.
- [5] Jimbo, M, Miwa, T. and Takeyama, Y., Counting minimal form factors of the restricted sine-Gordon model, math-ph/0303059.
- [6] Jimbo, M, Miwa, T., Mukhin, E. and Takeyama, Y., Form factors and action of $U_{\sqrt{-1}}(\tilde{sl_2})$ on infinite-cycles, *Comm. Math. Phys.* **245** (2004), 551-576. .
- [7] Melzer, E., The many faces of a character, *Lett. Math. Phys.* **31** (1994), 233–246.
- [8] Matsumura, H., Commutative Algebra, Benjamin, New York, 1970.
- [9] Nakayashiki, A., On the cohomologies of theta divisors of hyperelliptic Jacobians, *Contemporary Math.* **309** (2002), 177-183.
- [10] Nakayashiki, A., Residues of q -hypergeometric integrals and characters of affine Lie algebras, *Comm. Math. Phys.* **240** (2003), 197-241.
- [11] Nakayashiki, A., The chiral space of local operators in SU(2)-invariant Thirring model, *Comm. Math. Phys.* **245** (2004), 279-296. .
- [12] Nakayashiki, A. and Smirnov, F., Cohomologies of affine hyperelliptic Jacobi varieties and integrable systems, *Comm. Math. Phys.* **217** (2001), 623-652.
- [13] Nakayashiki, A. and Smirnov, F., Euler characteristics of theta divisors of Jacobians for spectral curves, *CRM Proc. and Lect. Notes* **32**, Vadim B. Kuznetsov, ed. (2002), 239-246,
- [14] Zamolodchikov, A. B., Integrable field theory from conformal field theory, *Adv. Stud. in Pure Math.* **19** (1989), 641–674.